


r (centimeters)	0	1	2	2.5	4
$f(r)$ (milligrams per square centimeter)	1	2	6	10	18

1. The density of a bacteria population in a circular petri dish at a distance r centimeters from the center of the dish is given by an increasing, differentiable function f , where $f(r)$ is measured in milligrams per square centimeter. Values of $f(r)$ for selected values of r are given in the table above.
- (a) Use the data in the table to estimate $f'(2.25)$. Using correct units, interpret the meaning of your answer in the context of this problem.
- (b) The total mass, in milligrams, of bacteria in the petri dish is given by the integral expression $2\pi \int_0^4 r f(r) dr$. Approximate the value of $2\pi \int_0^4 r f(r) dr$ using a right Riemann sum with the four subintervals indicated by the data in the table.
- (c) Is the approximation found in part (b) an overestimate or underestimate of the total mass of bacteria in the petri dish? Explain your reasoning.
- (d) The density of bacteria in the petri dish, for $1 \leq r \leq 4$, is modeled by the function g defined by $g(r) = 2 - 16(\cos(1.57\sqrt{r}))^3$. For what value of k , $1 < k < 4$, is $g(k)$ equal to the average value of $g(r)$ on the interval $1 \leq r \leq 4$?
2. For time $t \geq 0$, a particle moves in the xy -plane with position $(x(t), y(t))$ and velocity vector $\left\langle (t-1)e^{t^2}, \sin(t^{1.25}) \right\rangle$. At time $t = 0$, the position of the particle is $(-2, 5)$.
- (a) Find the speed of the particle at time $t = 1.2$. Find the acceleration vector of the particle at time $t = 1.2$.
- (b) Find the total distance traveled by the particle over the time interval $0 \leq t \leq 1.2$.
- (c) Find the coordinates of the point at which the particle is farthest to the left for $t \geq 0$. Explain why there is no point at which the particle is farthest to the right for $t \geq 0$.




3. A company designs spinning toys using the family of functions $y = cx\sqrt{4 - x^2}$, where c is a positive constant. The figure above shows the region in the first quadrant bounded by the x -axis and the graph of $y = cx\sqrt{4 - x^2}$, for some c . Each spinning toy is in the shape of the solid generated when such a region is revolved about the x -axis. Both x and y are measured in inches.

(a) Find the area of the region in the first quadrant bounded by the x -axis and the graph of $y = cx\sqrt{4 - x^2}$ for $c = 6$.

(b) It is known that, for $y = cx\sqrt{4 - x^2}$, $\frac{dy}{dx} = \frac{c(4 - 2x^2)}{\sqrt{4 - x^2}}$. For a particular spinning toy, the radius of the largest cross-sectional circular slice is 1.2 inches. What is the value of c for this spinning toy?

(c) For another spinning toy, the volume is 2π cubic inches. What is the value of c for this spinning toy?



Graph of f

4. Let f be a continuous function defined on the closed interval $-4 \leq x \leq 6$. The graph of f , consisting of four line segments, is shown above. Let G be the function defined by $G(x) = \int_0^x f(t) dt$.

(a) On what open intervals is the graph of G concave up? Give a reason for your answer.

(b) Let P be the function defined by $P(x) = G(x) \cdot f(x)$. Find $P'(3)$.

(c) Find $\lim_{x \rightarrow 2} \frac{G(x)}{x^2 - 2x}$.

(d) Find the average rate of change of G on the interval $[-4, 2]$. Does the Mean Value Theorem guarantee a value c , $-4 < c < 2$, for which $G'(c)$ is equal to this average rate of change? Justify your answer.

5. Let $y = f(x)$ be the particular solution to the differential equation $\frac{dy}{dx} = y \cdot (x \ln x)$ with initial condition $f(1) = 4$. It can be shown that $f''(1) = 4$.

(a) Write the second-degree Taylor polynomial for f about $x = 1$. Use the Taylor polynomial to approximate $f(2)$.

(b) Use Euler's method, starting at $x = 1$ with two steps of equal size, to approximate $f(2)$. Show the work that leads to your answer.

(c) Find the particular solution $y = f(x)$ to the differential equation $\frac{dy}{dx} = y \cdot (x \ln x)$ with initial condition $f(1) = 4$.

6. The function g has derivatives of all orders for all real numbers. The Maclaurin series for g is given by

$$g(x) = \sum_{n=0}^{\infty} \frac{(-1)^n x^n}{2e^n + 3} \text{ on its interval of convergence.}$$

(a) State the conditions necessary to use the integral test to determine convergence of the series $\sum_{n=0}^{\infty} \frac{1}{e^n}$.

Use the integral test to show that $\sum_{n=0}^{\infty} \frac{1}{e^n}$ converges.

(b) Use the limit comparison test with the series $\sum_{n=0}^{\infty} \frac{1}{e^n}$ to show that the series $g(1) = \sum_{n=0}^{\infty} \frac{(-1)^n}{2e^n + 3}$ converges absolutely.

(c) Determine the radius of convergence of the Maclaurin series for g .

(d) The first two terms of the series $g(1) = \sum_{n=0}^{\infty} \frac{(-1)^n}{2e^n + 3}$ are used to approximate $g(1)$. Use the alternating series error bound to determine an upper bound on the error of the approximation.

2021 BC Free Response Answers

- 1)
- a) $8 \text{ mg/cm}^2/\text{cm}$; At distance of 2.25 cm, $f'(2.25)$ is rate of change of density.
 - b) $269\pi = 845.088$
 - c) Overestimate because $rf(r)$ is increasing
 - d) 2.497
- 2)
- a) Speed = 1.271, accel. = $\langle 6.247, 0.405 \rangle$
 - b) 1.010
 - c) Furthest left at $(-2.604, 5.410)$, not furthest right because continues right for all $t > 1$
- 3)
- a) 16
 - b) 0.6
 - c) $\sqrt{\frac{15}{32}}$
- 4)
- a) $-4 < x < -2$ and $2 < x < 6$; f is increasing
 - b) 5.5
 - c) -2
 - d) $\frac{8}{3}$, yes because G is differentiable and continuous on the interval
- 5)
- a) $4 + 2(x - 1)^2$; 6
 - b) $4 + 3 \ln 1.5$
 - c) $y = e^{\left(\frac{x^2 \ln x}{2} - \frac{x^2}{4} + \ln 4 + \frac{1}{4}\right)}$
- 6)
- a) e^{-x} is positive, decreasing, and continuous
 - b)
 - c) e
 - d) $\frac{1}{2e^2 + 3}$