

Spring Break Assignment

ALL WORK MUST GO ON A SEPARATE SHEET OF PAPER. IF YOU DON'T KNOW HOW TO DO A PROBLEM, COPY THE ENTIRE PROBLEM AND ALL CHOICES ONTO YOUR PAPER.

No Calculators Allowed:

- At time $t \geq 0$, a particle moving in the xy -plane has velocity vector given by $v(t) = \langle t^2, 5t \rangle$. What is the acceleration vector of the particle at time $t = 3$?
(A) $\langle 9, \frac{45}{2} \rangle$ (B) $\langle 6, 5 \rangle$ (C) $\langle 2, 0 \rangle$ (D) $\sqrt{306}$ (E) $\sqrt{61}$
- $\int x e^{x^2} dx =$
(A) $\frac{1}{2} e^{x^2} + C$ (B) $e^{x^2} + C$ (C) $x e^{x^2} + C$ (D) $\frac{1}{2} e^{2x} + C$ (E) $e^{2x} + C$
- $\lim_{x \rightarrow 0} \frac{\sin x \cos x}{x}$ is
(A) -1 (B) 0 (C) 1 (D) $\frac{\pi}{4}$ (E) nonexistent
- Consider the series $\sum_{n=1}^{\infty} \frac{e^n}{n!}$. If the ratio test is applied to the series, which of the following inequalities results, implying that the series converges?
(A) $\lim_{n \rightarrow \infty} \frac{e}{n!} < 1$
(B) $\lim_{n \rightarrow \infty} \frac{n!}{e} < 1$
(C) $\lim_{n \rightarrow \infty} \frac{n+1}{e} < 1$
(D) $\lim_{n \rightarrow \infty} \frac{e}{n+1} < 1$
(E) $\lim_{n \rightarrow \infty} \frac{e}{(n+1)!} < 1$

5. Which of the following gives the length of the path described by the parametric equations $x = \sin(t^3)$ and $y = e^{5t}$ from $t = 0$ to $t = \pi$?

(A) $\int_0^\pi \sqrt{\sin^2(t^3) + e^{10t}} dt$

(B) $\int_0^\pi \sqrt{\cos^2(t^3) + e^{10t}} dt$

(C) $\int_0^\pi \sqrt{9t^4 \cos^2(t^3) + 25e^{10t}} dt$

(D) $\int_0^\pi \sqrt{3t^2 \cos(t^3) + 5e^{5t}} dt$

(E) $\int_0^\pi \sqrt{\cos^2(3t^2) + e^{10t}} dt$

$$f(x) = \begin{cases} \frac{x^2 - 4}{x - 2} & \text{if } x \neq 2 \\ 1 & \text{if } x = 2 \end{cases}$$

6. Let f be the function defined above. Which of the following statements about f are true?

I. f has a limit at $x = 2$.

II. f is continuous at $x = 2$.

III. f is differentiable at $x = 2$.

(A) I only

(B) II only

(C) III only

(D) I and II only

(E) I, II, and III

7. Given that $y(1) = -3$ and $\frac{dy}{dx} = 2x + y$, what is the approximation for $y(2)$ if Euler's method is used with a step size of 0.5, starting at $x = 1$?

(A) -5

(B) -4.25

(C) -4

(D) -3.75

(E) -3.5

x	2	3	5	8	13
$f(x)$	6	-2	-1	3	9

8. The function f is continuous on the closed interval $[2, 13]$ and has values as shown in the table above. Using the intervals $[2, 3]$, $[3, 5]$, $[5, 8]$, and $[8, 13]$, what is the approximation of $\int_2^{13} f(x) dx$ obtained from a left Riemann sum?


(A) 6

(B) 14

(C) 28

(D) 32

(E) 50



Graph of f

9. The graph of the piecewise linear function f is shown in the figure above. If $g(x) = \int_{-2}^x f(t) dt$, which of the following values is greatest?

- (A) $g(-3)$ (B) $g(-2)$ (C) $g(0)$ (D) $g(1)$ (E) $g(2)$

10. In the xy -plane, what is the slope of the line tangent to the graph of $x^2 + xy + y^2 = 7$ at the point $(2, 1)$?

- (A) $-\frac{4}{3}$ (B) $-\frac{5}{4}$ (C) -1 (D) $-\frac{4}{5}$ (E) $-\frac{3}{4}$

11. Let R be the region between the graph of $y = e^{-2x}$ and the x -axis for $x \geq 3$. The area of R is

- (A) $\frac{1}{2e^6}$ (B) $\frac{1}{e^6}$ (C) $\frac{2}{e^6}$ (D) $\frac{\pi}{2e^6}$ (E) infinite

12. Which of the following series converges for all real numbers x ?

(A) $\sum_{n=1}^{\infty} \frac{x^n}{n}$

(B) $\sum_{n=1}^{\infty} \frac{x^n}{n^2}$

(C) $\sum_{n=1}^{\infty} \frac{x^n}{\sqrt{n}}$

(D) $\sum_{n=1}^{\infty} \frac{e^n x^n}{n!}$

(E) $\sum_{n=1}^{\infty} \frac{n! x^n}{e^n}$

13. $\int_1^e \frac{x^2 + 1}{x} dx =$


- (A) $\frac{e^2 - 1}{2}$ (B) $\frac{e^2 + 1}{2}$ (C) $\frac{e^2 + 2}{2}$ (D) $\frac{e^2 - 1}{e^2}$ (E) $\frac{2e^2 - 8e + 6}{3e}$

x	0	1	2	3
$f''(x)$	5	0	-7	4

14. The polynomial function f has selected values of its second derivative f'' given in the table above. Which of the following statements must be true?
- (A) f is increasing on the interval $(0, 2)$.
 (B) f is decreasing on the interval $(0, 2)$.
 (C) f has a local maximum at $x = 1$.
 (D) The graph of f has a point of inflection at $x = 1$.
 (E) The graph of f changes concavity in the interval $(0, 2)$.
15. If $f(x) = (\ln x)^2$, then $f''(\sqrt{e}) =$
- (A) $\frac{1}{e}$ (B) $\frac{2}{e}$ (C) $\frac{1}{2\sqrt{e}}$ (D) $\frac{1}{\sqrt{e}}$ (E) $\frac{2}{\sqrt{e}}$
16. What are all values of x for which the series $\sum_{n=1}^{\infty} \left(\frac{2}{x^2 + 1}\right)^n$ converges?
- (A) $-1 < x < 1$
 (B) $x > 1$ only
 (C) $x \geq 1$ only
 (D) $x < -1$ and $x > 1$ only
 (E) $x \leq -1$ and $x \geq 1$
17. Let h be a differentiable function, and let f be the function defined by $f(x) = h(x^2 - 3)$. Which of the following is equal to $f'(2)$?
- (A) $h'(1)$ (B) $4h'(1)$ (C) $4h'(2)$ (D) $h'(4)$ (E) $4h'(4)$
18. In the xy -plane, the line $x + y = k$, where k is a constant, is tangent to the graph of $y = x^2 + 3x + 1$. What is the value of k ?
- (A) -3 (B) -2 (C) -1 (D) 0 (E) 1
19. $\int \frac{7x}{(2x-3)(x+2)} dx =$
- (A) $\frac{3}{2} \ln|2x-3| + 2 \ln|x+2| + C$
 (B) $3 \ln|2x-3| + 2 \ln|x+2| + C$
 (C) $3 \ln|2x-3| - 2 \ln|x+2| + C$
 (D) $-\frac{6}{(2x-3)^2} - \frac{2}{(x+2)^2} + C$
 (E) $-\frac{3}{(2x-3)^2} - \frac{2}{(x+2)^2} + C$

20. What is the sum of the series $1 + \ln 2 + \frac{(\ln 2)^2}{2!} + \dots + \frac{(\ln 2)^n}{n!} + \dots$?

- (A) $\ln 2$
- (B) $\ln(1 + \ln 2)$
- (C) 2
- (D) e^2
- (E) The series diverges.



21. A particle moves along a straight line. The graph of the particle's position $x(t)$ at time t is shown above for $0 < t < 6$. The graph has horizontal tangents at $t = 1$ and $t = 5$ and a point of inflection at $t = 2$. For what values of t is the velocity of the particle increasing?

- (A) $0 < t < 2$
- (B) $1 < t < 5$
- (C) $2 < t < 6$
- (D) $3 < t < 5$ only
- (E) $1 < t < 2$ and $5 < t < 6$

x	0	1
$f(x)$	2	4
$f'(x)$	6	-3
$g(x)$	-4	3
$g'(x)$	2	-1

22. The table above gives values of f , f' , g , and g' for selected values of x . If $\int_0^1 f'(x)g(x) dx = 5$, then

$$\int_0^1 f(x)g'(x) dx =$$

- (A) -14
- (B) -13
- (C) -2
- (D) 7
- (E) 15

23. If $f(x) = x \sin(2x)$, which of the following is the Taylor series for f about $x = 0$?


(A) $x - \frac{x^3}{2!} + \frac{x^5}{4!} - \frac{x^7}{6!} + \dots$

(B) $x - \frac{4x^3}{2!} + \frac{16x^5}{4!} - \frac{64x^7}{6!} + \dots$

(C) $2x - \frac{8x^3}{3!} + \frac{32x^5}{5!} - \frac{128x^7}{7!} + \dots$

(D) $2x^2 - \frac{2x^4}{3!} + \frac{2x^6}{5!} - \frac{2x^8}{7!} + \dots$

(E) $2x^2 - \frac{8x^4}{3!} + \frac{32x^6}{5!} - \frac{128x^8}{7!} + \dots$



24. Which of the following differential equations for a population P could model the logistic growth shown in the figure above?

(A) $\frac{dP}{dt} = 0.2P - 0.001P^2$

(B) $\frac{dP}{dt} = 0.1P - 0.001P^2$

(C) $\frac{dP}{dt} = 0.2P^2 - 0.001P$

(D) $\frac{dP}{dt} = 0.1P^2 - 0.001P$

(E) $\frac{dP}{dt} = 0.1P^2 + 0.001P$

$$f(x) = \begin{cases} cx + d & \text{for } x \leq 2 \\ x^2 - cx & \text{for } x > 2 \end{cases}$$

25. Let f be the function defined above, where c and d are constants. If f is differentiable at $x = 2$, what is the value of $c + d$?

- (A) -4 (B) -2 (C) 0 (D) 2 (E) 4

26. Which of the following expressions gives the total area enclosed by the polar curve $r = \sin^2 \theta$ shown in the figure above?


(A) $\frac{1}{2} \int_0^\pi \sin^2 \theta \, d\theta$

(B) $\int_0^\pi \sin^2 \theta \, d\theta$






(C) $\frac{1}{2} \int_0^\pi \sin^4 \theta \, d\theta$

(D) $\int_0^\pi \sin^4 \theta \, d\theta$

(E) $2 \int_0^\pi \sin^4 \theta \, d\theta$



27. Which of the following could be the slope field for the differential equation $\frac{dy}{dx} = y^2 - 1$?




28. In the xy -plane, a particle moves along the parabola $y = x^2 - x$ with a constant speed of $2\sqrt{10}$ units per second.

If $\frac{dx}{dt} > 0$, what is the value of $\frac{dy}{dt}$ when the particle is at the point $(2, 2)$?


- (A) $\frac{2}{3}$ (B) $\frac{2\sqrt{10}}{3}$ (C) 3 (D) 6 (E) $6\sqrt{10}$

Calculators Allowed:



76. The graph of f' , the derivative of a function f , is shown above. The domain of f is the open interval $0 < x < d$. Which of the following statements is true?

- (A) f has a local minimum at $x = c$.
(B) f has a local maximum at $x = b$.
(C) The graph of f has a point of inflection at $(a, f(a))$.
(D) The graph of f has a point of inflection at $(b, f(b))$.
(E) The graph of f is concave up on the open interval (c, d) .
77. Water is pumped out of a lake at the rate $R(t) = 12\sqrt{\frac{t}{t+1}}$ cubic meters per minute, where t is measured in minutes. How much water is pumped from time $t = 0$ to $t = 5$?
- (A) 9.439 cubic meters
(B) 10.954 cubic meters
(C) 43.816 cubic meters
(D) 47.193 cubic meters
(E) 54.772 cubic meters



78. The graph of a function f is shown above. For which of the following values of c does $\lim_{x \rightarrow c} f(x) = 1$?

- (A) 0 only
- (B) 0 and 3 only
- (C) -2 and 0 only
- (D) -2 and 3 only
- (E) -2, 0, and 3

79. Let f be a positive, continuous, decreasing function such that $a_n = f(n)$. If $\sum_{n=1}^{\infty} a_n$ converges to k , which of the following must be true?

- (A) $\lim_{n \rightarrow \infty} a_n = k$
- (B) $\int_1^n f(x) dx = k$
- (C) $\int_1^{\infty} f(x) dx$ diverges.
- (D) $\int_1^{\infty} f(x) dx$ converges.
- (E) $\int_1^{\infty} f(x) dx = k$

80. The derivative of the function f is given by $f'(x) = x^2 \cos(x^2)$. How many points of inflection does the graph of f have on the open interval $(-2, 2)$?

- (A) One
- (B) Two
- (C) Three
- (D) Four
- (E) Five

81. Let f and g be continuous functions for $a \leq x \leq b$. If $a < c < b$, $\int_a^b f(x) dx = P$, $\int_c^b f(x) dx = Q$,

$$\int_a^b g(x) dx = R, \text{ and } \int_c^b g(x) dx = S, \text{ then } \int_a^c (f(x) - g(x)) dx =$$

- (A) $P - Q + R - S$
- (B) $P - Q - R + S$
- (C) $P - Q - R - S$
- (D) $P + Q - R - S$
- (E) $P + Q - R + S$

82. If $\sum_{n=1}^{\infty} a_n$ diverges and $0 \leq a_n \leq b_n$ for all n , which of the following statements must be true?

(A) $\sum_{n=1}^{\infty} (-1)^n a_n$ converges.

(B) $\sum_{n=1}^{\infty} (-1)^n b_n$ converges.

(C) $\sum_{n=1}^{\infty} (-1)^n b_n$ diverges.

(D) $\sum_{n=1}^{\infty} b_n$ converges.

(E) $\sum_{n=1}^{\infty} b_n$ diverges.

83. What is the area enclosed by the curves $y = x^3 - 8x^2 + 18x - 5$ and $y = x + 5$?

- (A) 10.667 (B) 11.833 (C) 14.583 (D) 21.333 (E) 32

84. Let f be a function with $f(3) = 2$, $f'(3) = -1$, $f''(3) = 6$, and $f'''(3) = 12$. Which of the following is the third-degree Taylor polynomial for f about $x = 3$?

(A) $2 - (x - 3) + 3(x - 3)^2 + 2(x - 3)^3$

(B) $2 - (x - 3) + 3(x - 3)^2 + 4(x - 3)^3$

(C) $2 - (x - 3) + 6(x - 3)^2 + 12(x - 3)^3$

(D) $2 - x + 3x^2 + 2x^3$

(E) $2 - x + 6x^2 + 12x^3$

85. A particle moves on the x -axis with velocity given by $v(t) = 3t^4 - 11t^2 + 9t - 2$ for $-3 \leq t \leq 3$. How many times does the particle change direction as t increases from -3 to 3 ?

- (A) Zero (B) One (C) Two (D) Three (E) Four

86. On the graph of $y = f(x)$, the slope at any point (x, y) is twice the value of x . If $f(2) = 3$, what is the value of $f(3)$?

- (A) 6 (B) 7 (C) 8 (D) 9 (E) 10

87. An object traveling in a straight line has position $x(t)$ at time t . If the initial position is $x(0) = 2$ and the velocity of the object is $v(t) = \sqrt[3]{1 + t^2}$, what is the position of the object at time $t = 3$?

- (A) 0.431 (B) 2.154 (C) 4.512 (D) 6.512 (E) 17.408

88. For all values of x , the continuous function f is positive and decreasing. Let g be the function given by

$$g(x) = \int_2^x f(t) dt. \text{ Which of the following could be a table of values for } g?$$

(A)

x	$g(x)$
1	-2
2	0
3	1

(B)

x	$g(x)$
1	-2
2	0
3	3

(C)

x	$g(x)$
1	1
2	0
3	-2

(D)

x	$g(x)$
1	2
2	0
3	-1

(E)

x	$g(x)$
1	3
2	0
3	2

89. The function f is continuous for $-2 \leq x \leq 2$ and $f(-2) = f(2) = 0$. If there is no c , where $-2 < c < 2$, for which $f'(c) = 0$, which of the following statements must be true?

- (A) For $-2 < k < 2$, $f'(k) > 0$.
- (B) For $-2 < k < 2$, $f'(k) < 0$.
- (C) For $-2 < k < 2$, $f'(k)$ exists.
- (D) For $-2 < k < 2$, $f'(k)$ exists, but f' is not continuous.
- (E) For some k , where $-2 < k < 2$, $f'(k)$ does not exist.


x	$f(x)$	$g(x)$	$f'(x)$	$g'(x)$
-1	-5	1	3	0
0	-2	0	1	1
1	0	-3	0	0.5
2	5	-1	5	2

90. The table above gives values of the differentiable functions f and g and of their derivatives f' and g' , at selected values of x . If $h(x) = f(g(x))$, what is the slope of the graph of h at $x = 2$?

- (A) -10 (B) -6 (C) 5 (D) 6 (E) 10

91. Let f be the function given by $f(x) = \int_{1/3}^x \cos\left(\frac{1}{t^2}\right) dt$ for $\frac{1}{3} \leq x \leq 1$. At which of the following values of x does f attain a relative maximum?


- (A) 0.357 and 0.798 (B) 0.4 and 0.564 (C) 0.4 only (D) 0.461 (E) 0.999



92. The figure above shows the graphs of the functions f and g . The graphs of the lines tangent to the graph of g at $x = -3$ and $x = 1$ are also shown. If $B(x) = g(f(x))$, what is $B'(-3)$?

- (A) $-\frac{1}{2}$ (B) $-\frac{1}{6}$ (C) $\frac{1}{6}$ (D) $\frac{1}{3}$ (E) $\frac{1}{2}$

Calculators Allowed:



1. Let R be the region bounded by the graphs of $y = \sin(\pi x)$ and $y = x^3 - 4x$, as shown in the figure above.
 - (a) Find the area of R .
 - (b) The horizontal line $y = -2$ splits the region R into two parts. Write, but do not evaluate, an integral expression for the area of the part of R that is below this horizontal line.
 - (c) The region R is the base of a solid. For this solid, each cross section perpendicular to the x -axis is a square. Find the volume of this solid.
 - (d) The region R models the surface of a small pond. At all points in R at a distance x from the y -axis, the depth of the water is given by $h(x) = 3 - x$. Find the volume of water in the pond.


t (hours)	0	1	3	4	7	8	9
$L(t)$ (people)	120	156	176	126	150	80	0

2. Concert tickets went on sale at noon ($t = 0$) and were sold out within 9 hours. The number of people waiting in line to purchase tickets at time t is modeled by a twice-differentiable function L for $0 \leq t \leq 9$. Values of $L(t)$ at various times t are shown in the table above.
- Use the data in the table to estimate the rate at which the number of people waiting in line was changing at 5:30 P.M. ($t = 5.5$). Show the computations that lead to your answer. Indicate units of measure.
 - Use a trapezoidal sum with three subintervals to estimate the average number of people waiting in line during the first 4 hours that tickets were on sale.
 - For $0 \leq t \leq 9$, what is the fewest number of times at which $L'(t)$ must equal 0? Give a reason for your answer.
 - The rate at which tickets were sold for $0 \leq t \leq 9$ is modeled by $r(t) = 550te^{-t/2}$ tickets per hour. Based on the model, how many tickets were sold by 3 P.M. ($t = 3$), to the nearest whole number?

x	$h(x)$	$h'(x)$	$h''(x)$	$h'''(x)$	$h^{(4)}(x)$
1	11	30	42	99	18
2	80	128	$\frac{488}{3}$	$\frac{448}{3}$	$\frac{584}{9}$
3	317	$\frac{753}{2}$	$\frac{1383}{4}$	$\frac{3483}{16}$	$\frac{1125}{16}$

3. Let h be a function having derivatives of all orders for $x > 0$. Selected values of h and its first four derivatives are indicated in the table above. The function h and these four derivatives are increasing on the interval $1 \leq x \leq 3$.
- Write the first-degree Taylor polynomial for h about $x = 2$ and use it to approximate $h(1.9)$. Is this approximation greater than or less than $h(1.9)$? Explain your reasoning.
 - Write the third-degree Taylor polynomial for h about $x = 2$ and use it to approximate $h(1.9)$.
 - Use the Lagrange error bound to show that the third-degree Taylor polynomial for h about $x = 2$ approximates $h(1.9)$ with error less than 3×10^{-4} .

No Calculators Allowed:




Graph of v

4. A particle moves along the x -axis so that its velocity at time t , for $0 \leq t \leq 6$, is given by a differentiable function v whose graph is shown above. The velocity is 0 at $t = 0$, $t = 3$, and $t = 5$, and the graph has horizontal tangents at $t = 1$ and $t = 4$. The areas of the regions bounded by the t -axis and the graph of v on the intervals $[0, 3]$, $[3, 5]$, and $[5, 6]$ are 8, 3, and 2, respectively. At time $t = 0$, the particle is at $x = -2$.
- For $0 \leq t \leq 6$, find both the time and the position of the particle when the particle is farthest to the left. Justify your answer.
 - For how many values of t , where $0 \leq t \leq 6$, is the particle at $x = -8$? Explain your reasoning.
 - On the interval $2 < t < 3$, is the speed of the particle increasing or decreasing? Give a reason for your answer.
 - During what time intervals, if any, is the acceleration of the particle negative? Justify your answer.

-
5. The derivative of a function f is given by $f'(x) = (x - 3)e^x$ for $x > 0$, and $f(1) = 7$.
- The function f has a critical point at $x = 3$. At this point, does f have a relative minimum, a relative maximum, or neither? Justify your answer.
 - On what intervals, if any, is the graph of f both decreasing and concave up? Explain your reasoning.
 - Find the value of $f(3)$.

6. Consider the logistic differential equation $\frac{dy}{dt} = \frac{y}{8}(6 - y)$. Let $y = f(t)$ be the particular solution to the differential equation with $f(0) = 8$.

- A slope field for this differential equation is given below. Sketch possible solution curves through the points $(3, 2)$ and $(0, 8)$.



- Use Euler's method, starting at $t = 0$ with two steps of equal size, to approximate $f(1)$.
- Write the second-degree Taylor polynomial for f about $t = 0$, and use it to approximate $f(1)$.
- What is the range of f for $t \geq 0$?

Spring Break Assignment

- | | | | | | | | | | |
|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| 1. B | 2. A | 3. C | 4. D | 5. C | 6. A | 7. D | 8. B | 9. D | 10. B |
| 11. A | 12. D | 13. B | 14. E | 15. A | 16. D | 17. B | 18. A | 19. A | 20. C |
| 21. A | 22. E | 23. E | 24. A | 25. B | 26. D | 27. A | 28. D | 76. D | 77. D |
| 78. C | 79. D | 80. E | 81. B | 82. E | 83. B | 84. A | 85. C | 86. C | 87. D |
| 88. A | 89. E | 90. D | 91. D | 92. B | | | | | |

1.

(a) $\sin(\pi x) = x^3 - 4x$ at $x = 0$ and $x = 2$

$$\text{Area} = \int_0^2 (\sin(\pi x) - (x^3 - 4x)) dx = 4$$

(b) $x^3 - 4x = -2$ at $r = 0.5391889$ and $s = 1.6751309$

$$\text{The area of the stated region is } \int_r^s (-2 - (x^3 - 4x)) dx$$

(c) $\text{Volume} = \int_0^2 (\sin(\pi x) - (x^3 - 4x))^2 dx = 9.978$

(d) $\text{Volume} = \int_0^2 (3 - x)(\sin(\pi x) - (x^3 - 4x)) dx = 8.369$ or 8.370

2.

(a) $L'(5.5) \approx \frac{L(7) - L(4)}{7 - 4} = \frac{150 - 126}{3} = 8$ people per hour

(b) The average number of people waiting in line during the first 4 hours is approximately

$$\frac{1}{4} \left(\frac{L(0) + L(1)}{2} (1 - 0) + \frac{L(1) + L(3)}{2} (3 - 1) + \frac{L(3) + L(4)}{2} (4 - 3) \right) = 155.25 \text{ people}$$

(c) L is differentiable on $[0, 9]$ so the Mean Value Theorem implies $L'(t) > 0$ for some t in $(1, 3)$ and some t in $(4, 7)$. Similarly, $L'(t) < 0$ for some t in $(3, 4)$ and some t in $(7, 8)$. Then, since L' is continuous on $[0, 9]$, the Intermediate Value Theorem implies that $L'(t) = 0$ for at least three values of t in $[0, 9]$.

OR

The continuity of L on $[1, 4]$ implies that L attains a maximum value there. Since $L(3) > L(1)$ and $L(3) > L(4)$, this maximum occurs on $(1, 4)$. Similarly, L attains a minimum on $(3, 7)$ and a maximum on $(4, 8)$. L is differentiable, so $L'(t) = 0$ at each relative extreme point on $(0, 9)$. Therefore $L'(t) = 0$ for at least three values of t in $[0, 9]$.

[Note: There is a function L that satisfies the given conditions with $L'(t) = 0$ for exactly three values of t .]

(d) $\int_0^3 r(t) dt = 972.784$

There were approximately 973 tickets sold by 3 P.M.

3.

(a) $P_1(x) = 80 + 128(x - 2)$, so $h(1.9) \approx P_1(1.9) = 67.2$

$P_1(1.9) < h(1.9)$ since h' is increasing on the interval $1 \leq x \leq 3$.

(b) $P_3(x) = 80 + 128(x - 2) + \frac{488}{6}(x - 2)^2 + \frac{448}{18}(x - 2)^3$

$h(1.9) \approx P_3(1.9) = 67.988$

(c) The fourth derivative of h is increasing on the interval

$1 \leq x \leq 3$, so $\max_{1.9 \leq x \leq 2} |h^{(4)}(x)| = \frac{584}{9}$.

Therefore, $|h(1.9) - P_3(1.9)| \leq \frac{584}{9} \frac{|1.9 - 2|^4}{4!}$
 $= 2.7037 \times 10^{-4}$
 $< 3 \times 10^{-4}$

5.

(a) $f'(x) < 0$ for $0 < x < 3$ and $f'(x) > 0$ for $x > 3$

Therefore, f has a relative minimum at $x = 3$.

(b) $f''(x) = e^x + (x - 3)e^x = (x - 2)e^x$
 $f''(x) > 0$ for $x > 2$

$f'(x) < 0$ for $0 < x < 3$

Therefore, the graph of f is both decreasing and concave up on the interval $2 < x < 3$.

(c) $f(3) = f(1) + \int_1^3 f'(x) dx = 7 + \int_1^3 (x - 3)e^x dx$

$u = x - 3 \quad dv = e^x dx$
 $du = dx \quad v = e^x$

$f(3) = 7 + (x - 3)e^x \Big|_1^3 - \int_1^3 e^x dx$
 $= 7 + ((x - 3)e^x - e^x) \Big|_1^3$
 $= 7 + 3e - e^3$

4.

(a) Since $v(t) < 0$ for $0 < t < 3$ and $5 < t < 6$, and $v(t) > 0$ for $3 < t < 5$, we consider $t = 3$ and $t = 6$.

$x(3) = -2 + \int_0^3 v(t) dt = -2 - 8 = -10$

$x(6) = -2 + \int_0^6 v(t) dt = -2 - 8 + 3 - 2 = -9$

Therefore, the particle is farthest left at time $t = 3$ when its position is $x(3) = -10$.

(b) The particle moves continuously and monotonically from $x(0) = -2$ to $x(3) = -10$. Similarly, the particle moves continuously and monotonically from $x(3) = -10$ to $x(5) = -7$ and also from $x(5) = -7$ to $x(6) = -9$.


By the Intermediate Value Theorem, there are three values of t for which the particle is at $x(t) = -8$.

(c) The speed is decreasing on the interval $2 < t < 3$ since on this interval $v < 0$ and v is increasing.

(d) The acceleration is negative on the intervals $0 < t < 1$ and $4 < t < 6$ since velocity is decreasing on these intervals.

6.

(a)



(b) $f\left(\frac{1}{2}\right) \approx 8 + (-2)\left(\frac{1}{2}\right) = 7$

$f(1) \approx 7 + \left(-\frac{7}{8}\right)\left(\frac{1}{2}\right) = \frac{105}{16}$

(c) $\frac{d^2y}{dt^2} = \frac{1}{8} \frac{dy}{dt} (6 - y) + \frac{y}{8} \left(-\frac{dy}{dt}\right)$

$f(0) = 8; f'(0) = \frac{dy}{dt} \Big|_{t=0} = \frac{8}{8}(6 - 8) = -2; \text{ and}$

$f''(0) = \frac{d^2y}{dt^2} \Big|_{t=0} = \frac{1}{8}(-2)(-2) + \frac{8}{8}(2) = \frac{5}{2}$

The second-degree Taylor polynomial for f about

$t = 0$ is $P_2(t) = 8 - 2t + \frac{5}{4}t^2$.

$f(1) \approx P_2(1) = \frac{29}{4}$

(d) The range of f for $t \geq 0$ is $6 < y \leq 8$.